

Comparing the Properties of ICME-Induced Forbush Decreases at Earth and Mars Johan L. Freiherr von Forstner¹ Jingnan Guo^{1,2} Robert F. Wimmer-Schweingruber¹ Mateja Dumbović^{3,4} Miho Janvier⁵

Pascal Démoulin⁶ Astrid Veronig⁴ Manuela Temmer⁴ Athanasios Papaioannou⁷ Sergio Dasso⁸ Donald M. Hassler⁹ Cary J. Zeitlin¹⁰

Introduction

Forbush decreases (FDs) are short-term drops in the flux of galactic cosmic rays (GCR) caused by the shielding from magnetic structures in the solar wind. They can be related to **interplanetary coronal mass** ejections (ICMEs) or corotating interaction regions (CIRs). FDs can be observed at Earth, e.g. using neutron monitors, but also at other locations in the solar system, such as on the surface of Mars with the Radiation Assessment Detector (RAD) instrument onboard Mars Science Laboratory (MSL).

FDs are often used as a proxy for detecting the arrival of ICMEs or CIRs, especially when sufficient in situ solar wind measurements are not available. But with sufficient understanding of the FD physics, it is also possible to gain more information about the heliospheric structures than just their arrival times. We demonstrate such an approach, focusing on ICME-related FDs observed at Earth and Mars.



Definitions

 y/y_{max} : FD profile (y(t)) normalized to the onset value

 Δy : Drop ratio of the FD (in %)

 Δt : Duration of the FD decrease phase (in h)

 m_{max} : Maximum decrease rate during the decrease phase (in % h⁻¹) S, E Sheath and ejecta regions of the ICME

Observations I

Using our catalog of 45 ICME-induced FDs observed at Mars [2], we find a **linear correlation** of Δy and m_{max} (Figure 1). This correlation was already found at Earth [3, 4] and is confirmed by the subset of 14 events also seen on Earth (South Pole neutron monitor). However, the linear regression $\Delta y = A \cdot m_{\max} + B$ at Mars has a steeper slope A than at Earth by a factor of $E = A_{\text{Mars}}/A_{\text{Earth}} = 1.9 \pm 0.4$.



Figure 1. Correlation of Δy and m_{max} at Mars (orange) and Earth (blue). Orange points with blue outline mark the subset of events at Mars also seen at Earth.

¹University of Kiel ²University of Science and Technology of China, Hefei ⁶LESIA, Observatoire de Paris ⁷IAASARS, National Observatory of Athens

³University of Zagreb

Observations II

A similar trend is also visible in the larger catalog from Papaioannou et al. [5]. In this case, we applied a filter to only look at FDs associated with ICMEs (middle panel of Figure 2) as well as a threshold condition (1)

 $\Delta y \geq f \cdot \text{median}(\Delta y)$

to exclude FDs with low amplitude where the uncertainty of the Δy and $m_{\rm max}$ parameters can be large.



Figure 2. Same correlation plot for Earth and Mars based on a larger catalog from [5]. Events are plotted all together (top row) and separately for ICME-induced Forbush decreases and others (mainly CIRs). The threshold condition from Equation 1 is applied with f = 1.0.

The uncertainties of the fit parameters were calculated using a bootstrap method. An investigation of the dependence of A on the threshold parameter f shows that A_{Earth} and A_{Mars} become significantly different above $f \gtrsim 0.8$. At f = 1.0 (shown in Figure 2), the ratio E is 1.5 ± 0.2 .

Interpretation

Two possible reasons:

- Lower GCR energy observed at Mars Mars (MSL/RAD dose rate): mainly 1 GeV to 3 GeV protons Earth (global survey method): $10 \,\text{GV}$ rigidity $= 9.1 \,\text{GeV}$ protons \Rightarrow Only affects the FD magnitude, not duration.
- ICME evolution between Earth and Mars, e.g. expansion \Rightarrow Affects both FD magnitude and duration.

⁴University of Graz⁵Institut d'Astrophysique Spatiale, Orsay ⁸University of Buenos Aires⁹Southwest Research Institute, Boulder¹⁰Leidos, Houston

Modeling of m_{max} vs. Δy relation

Figure 3 shows an idealized schematic illustration of FDs at Earth and Mars, taking into account both effects mentioned above. It can be seen that only the expansion of the ICME structure results in a **steeper slope** A at Mars, while a change in GCR energy does not affect the slope.



b) Mars (expanded ICME, same GCR energy E_0)



c) Mars (expanded ICME, lower GCR energy E_1)



Figure 3. Cartoon illustration of the Δy vs m_{max} correlation for Earth and Mars.

We have performed analytical modeling calculations using PDB [6] and ForbMod [7], confirming that A is proportional to the duration of the ICME passage and does not depend on the GCR energy for both the sheath and ejecta regions of the ICME. As m_{max} usually occurs in the sheath, we suspect that our results are **mostly affected by the sheath region**. This was also validated by applying another threshold condition g to the catalog to only include FDs where m_{max} occurs close to the onset time, resulting in similar values of $E = 1.5 \pm 0.4$.

Thus, E should correspond to the **sheath broadening factor** between Earth and Mars.

Calculation of sheath broadening

As a comparison to the ratio obtained from FD measurements, we calculate the increase in sheath width S between 1 au and 1.5 au. In this region, it is probably dominated by two processes:

Pileup of ambient solar wind in front of the shock:

 $\Delta S_{\text{pileup}} \propto (v_{\text{shock}} - v_{\text{sw}}) \Delta t$

Expansion/contraction corresponding to the sheath velocity profile:

 $\Delta S_{\text{exp}} = (v_{\text{sheath,front}} - v_{\text{sheath,rear}})\Delta t$

Other effects are considered to be insignificant beyond 1 au for a firstorder approximation. Based on these equations and typical values obtained from statistical studies [8], we estimate the broadening factor $E = S_{\text{Mars}}/S_{\text{Earth}}$ to be between 1.2 and 1.8, comparable to the ratios of our linear regression slopes in the FD analysis.



Christian-Albrechts-Universität zu Kiel Institut für

Experimentelle und Angewandte Physik

Conclusion

We have estimated the factor E by which the size of an ICME sheath region increases between Earth and Mars based on analysis of only Forbush decrease measurements. This was made possible by the choice of two characteristic FD parameters which are linearly correlated, and the analytical modeling of the dependence of this relation on the properties of the ICME. Based on different data sets, we could calculate three results for E, which are in good agreement with each other (Table 1). Theoretical estimations of this factor also give similar results.

E value Description

 1.9 ± 0.4 FDs in Freiherr von Forstner et al. [2] catalog 1.5 ± 0.2 FDs in Papaioannou et al. [5] catalog, threshold f = 1 1.5 ± 0.4 FDs in Papaioannou et al. [5] catalog, threshold g = 0.31.2 to 1.8 Theoretical estimation of broadening factor

Table 1. Results for the sheath broadening factor E estimated in different parts of this study.

Our results for E extend on previous direct measurements of the sheath evolution closer to the Sun, such as by Janvier et al. [9]. They showed that the speed at which the broadening occurs decreases further away from the Sun, and our results between Earth and Mars agree with this trend.

References

Main citation: [1] J. L. Freiherr von Forstner et al. "Comparing the Properties of ICME-Induced Forbush decreases at Earth and Mars". In: JGR Space (submitted)

- [2] J. L. Freiherr von Forstner et al. "Tracking and Validating ICMEs Propagating Toward Mars Using STEREO Heliospheric Imagers Combined With Forbush Decreases Detected by MSL/RAD". In: Space Weather 17.4 (2019).
- [3] A. V. Belov. "Forbush effects and their connection with solar, interplanetary and geomagnetic phenomena". In: Proc. IAU 4.S257 (2008).
- [4] A. A. Abunin et al. "Forbush effects with a sudden and gradual onset". In: Ann. Geophys. 52.3 (2012).
- [5] A. Papaioannou et al. "A Catalogue of Forbush Decreases Recorded on the Surface of Mars from 2012 Until 2016: Comparison with Terrestrial FDs". In: Sol. Phys. 294.6 (2019).
- [6] G. Wibberenz et al. "Transient Effects and Disturbed Conditions". In: *Space Sci. Rev.* 83.1 (1998). [7] M. Dumbović et al. "An Analytical Diffusion–Expansion Model for Forbush Decreases Caused by Flux Ropes". In: ApJ 860.1 (2018).
- [8] J. J. Masías-Meza et al. "Superposed epoch study of ICME sub-structures near Earth and their effects on Galactic cosmic rays". In: A&A 592 (2016).
- M. Janvier et al. "Generic Magnetic Field Intensity Profiles of Interplanetary Coronal Mass Ejections at Mercury, Venus, and Earth From Superposed Epoch Analyses". In: JGR Space 124.2

Acknowledgements

RAD is supported by NASA (HEOMD) under JPL subcontract 1273039 to Southwest Research Institute and in Germany by DLR and DLR's Space Administration grants 50QM0501, 50QM1201, and 50QM1701 to the Christian Albrechts University, Kiel. We acknowledge the NMDB database (www.nmdb.eu), funded under the European Union's FP7 Programme (contract 213007), for providing data. The data from South Pole neutron monitor is provided by the University of Delaware

with support from the U.S. National Science Foundation under grant ANT-0838839. J.G. is supported by the Key Research Program of the Chinese Academy of Sciences (Grant No. XDPB11 and QYZDB-SSW-DQC015) and acknowledges LabEx P2iO at Université Paris-Saclay under the visiting researcher scheme "Emilie du Châtelet". The visit of J.G. to Paris observatory was funded by the LabEx Plas@Par, which is driven by Sorbonne Université. M.D. acknowledges partial funding from the EU H2020 MSCA grant agreements No. 745782 (ForbMod) and No. 824135 (SOLARNET) and support by the Croatian Science Foundation under the project 7549 (MSOC). A.P. would like to acknowledge the TRACER project funded by the National Observatory of Athens (NOA) (Project ID: 5063).

Scan the QR code to download this poster and find links to my

